

Chiral Symmetry of Baryons

Daisuke JIDO^{1,*} Makoto OKA^{2,**} and Atsushi HOSAKA^{3,***}

¹ *Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia-CSIC, Institutos de Investigación de Peterna, Aptdo. Correos 2085, 46071, Valencia, Spain*

² *Department of Physics, Tokyo Institute of Technology, Meguro, Tokyo 152-8551 Japan*

³ *Research Center for Nuclear Physics, Osaka University, Ibaraki, 567-0047 Japan*

(Received)

We study chiral symmetry aspects of the positive and negative parity baryons by identifying them with linear representations of the chiral group $SU(N_f) \otimes SU(N_f)$. It is shown that there are two distinctive schemes: naive and mirror assignments. We construct linear sigma models for baryons in the two assignments and examine their physical implications. Then we investigate properties of the naive and mirror nucleons microscopically by using QCD interpolating fields. Finally, we propose experiments to distinguish the two chiral assignments for the nucleon.

§1. Introduction

Chiral symmetry plays a unique role in hadron physics. In QCD, it is realized as the global flavor symmetry for the right and left handed quarks in the light flavor sector. The associated chiral group is then denoted as $SU(N_f)_R \otimes SU(N_f)_L$, where N_f is the number of light flavors. At low energies, however, it is spontaneously broken and only the diagonal vector symmetry remains to manifest,

$$SU(N_f)_R \otimes SU(N_f)_L \rightarrow SU(N_f)_V. \quad (1.1)$$

The spontaneous breakdown is accompanied by the appearance of the massless pseudo-scalar mesons, the pions and kaons, as the Nambu-Goldstone bosons^{1), 2)}. In the real world the mesons acquire finite but small masses due to small current quark masses. Then the pions play the central role in strong interaction physics at low energies. Chiral symmetry has been proved to be useful in understanding dynamics of low energy pions and nucleons as shown in the success of the current algebra based on conserved vector current (CVC) and partially conserved axial-vector current (PCAC)³⁾.

In spite of the success of such symmetry guided approaches, there still remains a fundamental question; what representations of the chiral group manifest in observed particles. In fact, such a question is not new. Prior to QCD, much study was performed in the late 60th^{4), 5)}. The reasons that we are now interested in this long

*) e-mail address: Daisuke.Jido@hal.ific.uv.es

**) e-mail address: oka@th-physics.titech.ac.jp

***) e-mail address: hosaka@rcnp.osaka-u.ac.jp

standing problem are: (1) Much progress has been made in the fundamental theory of QCD. (2) Relevant physics can now be accessible by experiments at the present day. (3) The problem of parity doubling, in particular, of the baryon sector has not been considered much. In this paper, we would like to focus on the last issue, the chiral symmetry of baryons.

To start with, let us recall the analysis by Weinberg about chiral representations for hadrons, in particular, for baryons⁴⁾. In a naive consideration of the relativistic nucleon field, it is natural to identify it with the fundamental representations of the chiral group $SU(2)_R \otimes SU(2)_L$ ^{*)}

$$\psi \sim (\tfrac{1}{2}, 0) \oplus (0, \tfrac{1}{2}). \quad (1.2)$$

Such a linear field may be coupled in a chiral invariant way with linear chiral mesons belonging to a representation $(\tfrac{1}{2}, \tfrac{1}{2}) \sim (\vec{\pi}, \sigma)$. This includes not only the pion but also the sigma meson σ , the existence of which is still under controversy. Then Weinberg motivated to eliminate such poorly established particles and constructed a theory only by well established particles, the pions, which are linear representations of the residual vector symmetry group⁶⁾. The elimination is made by imposing a constraint among the linear chiral fields, which necessarily lead to the non-linear realization of chiral symmetry. In the non-linear realization the lagrangian formulation serves a derivative expansion for amplitudes and has become the basis of the chiral perturbation theory⁷⁾⁻⁹⁾. In the non-linear sigma model, since the spontaneous breaking is assumed from the beginning, it does not seem easy to discuss dynamical issues such as the phase transition of chiral symmetry.

In this respect, linear sigma models are rather suited to the study of dynamical aspects of chiral symmetry¹⁰⁾. For instance, it can describe the spontaneous breaking within the model. Also, linear representations introduce chiral partners having different parities. This provides a systematic view on hadrons from chiral symmetry. For instance, mass splittings of positive and negative parity hadrons are generated by the spontaneous breaking of chiral symmetry and are governed by a unique mass scale. Furthermore, they become degenerate when chiral symmetry is restored. These seem plausible since observed mass differences of positive and negative parity hadrons are about 500 MeV independent of particle channels as shown in Fig. 1.

Now in QCD, the quark field q belongs to a linear representation of the chiral group as in Eq. (1.2). Then the chiral mesons are naively formed as quark bilinears:

$$(i\bar{q}\gamma_5\vec{\tau}q, \bar{q}q) \sim (\vec{\pi}, \sigma) \sim (\tfrac{1}{2}, \tfrac{1}{2}). \quad (1.3)$$

We observe that the scalar σ and pseudoscalar $\vec{\pi}$ are chiral partners among themselves. Similarly, we can construct the $(1, 0) \oplus (0, 1)$ representation for the vector (ρ) and axial-vector (a_1) mesons.

Then what is the chiral representation of the nucleon? Or, what is the chiral partner of the ground state nucleon? This is not a trivial question, since the nucleon is a composite of three quarks and is subject to strong interactions with the pions.

^{*)} Isospin values are used in order to denote representations of the isospin $SU(2)$ group. See, section 2 for definitions.




Fig. 1. Mass splittings of positive and negative parity hadrons in various channels. Data are taken from the Particle Data Booklet¹¹⁾. The uncertain mass of sigma (σ) is hatched.

The simplest possibility has been already given in (1.2); four components of the nucleon form a chiral representation among themselves, where it is not necessary to introduce negative parity nucleons as a chiral partner.

We can now list the relevant questions as follows:

- Is it possible to consider larger representations which include both positive and negative parity nucleons and relate them by chiral symmetry?
- If so, what are possible representations and what dynamical consequences are expected?
- What is the quark (QCD) origin of such realizations?
- Are there any experimental approaches to inform the chiral symmetry of baryons?

These are the topics we would like to study in this article.

Much of the material treated here is based on our former and currently progressing works¹²⁾⁻¹⁷⁾. In fact, stimulated by the similar motivations, Lee¹⁸⁾ and later DeTar and Kunihiro¹⁹⁾ previously constructed a theory in which positive and negative parity nucleons behave as members of a larger chiral representation. What we will explore here is then to clarify the meaning of their work from a fundamental point of view. A somewhat unusual way of introducing the two nucleons in their original works is now reformulated using group theoretical languages.

In section 2, after a basic introduction for chiral symmetry and its representations, we discuss group theoretical aspects of the two chiral representations for positive and negative parity nucleons. In section 3, we study linear sigma models in the two assignments and discuss phenomenological consequences. Several comments are made for theoretical aspects of the models. A microscopic origin of these two realizations is discussed in section 4, where various baryon interpolating fields are investigated. Since at the present time the theoretical understanding has reached only to a limited extent, we propose experiments which will help discussing the two assignments. For this, in section 5, we calculate reaction cross sections for the pion and eta productions at the threshold region. Final section is devoted to the summary of the present work.

§2. Representations of the Chiral Group for Baryons

Here we discuss linear representations of the chiral symmetry group for baryons starting from chiral symmetry for quarks. Finally we shall reach two possibilities to assign the chiral symmetry to two kinds of baryons, which makes different consequences of phenomenological properties of baryons.

2.1. Chiral Symmetry in QCD

QCD Lagrangian is given in a very simple form:

$$\mathcal{L}_{\text{QCD}} = \sum_f \bar{q}_f (i \not{D} - m_f) q_f - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}, \quad (2.1)$$

where $q_f (f = u, d, s, c, b, t)$ are fields of quarks with mass m_f , $F_{\mu\nu}^a$ denotes the field tensor of gluons with their color index a . For the light quarks ($f = u, d$ and s), there is an approximate symmetry independent of the chirality of the quarks, that is called chiral symmetry. The definition of chiral symmetry with N_f light flavor is given by flavor transformations on the right and left handed components (of the Lorentz group):

$$q_r \rightarrow e^{i\vec{t} \cdot \vec{r}} \psi_r, \quad q_l \rightarrow e^{i\vec{t} \cdot \vec{l}} \psi_l, \quad (2.2)$$

where $\vec{t}$ is a vector of $N_f^2 - 1$ generators of $SU(N_f)$, and $\vec{r}$ (and $\vec{l}$) are $N_f^2 - 1$ transformation parameters. The right and left handed components are projected out from a four component quark field q by

$$q = q_r + q_l, \quad (2.3)$$

$$q_r = \frac{1 + \gamma_5}{2} q, \quad q_l = \frac{1 - \gamma_5}{2} q. \quad (2.4)$$

The generators satisfy the following commutation relations and normalization:

$$[t^a, t^b] = i f_{abc} t^c, \quad \text{tr} (t^a t^b) = \frac{1}{2} \delta_{ab}, \quad (2.5)$$

where f_{abc} are the structure constants of $SU(N_f)$.

In the following discussions, we deal mostly with the case $N_f = 2$ for isospin $SU(2)$ symmetry. Thus the generators are given by

$$t^a = \frac{\tau^a}{2}, \quad (2.6)$$

where τ^a are the Pauli matrices, and the structure constants are $f_{abc} = \epsilon_{abc}$.

In the quantum theory, transformations of the field q are generated by the charge operators. For instance, the right handed field is transformed by the right chiral charges Q_R^a ,

$$q_r \rightarrow g_R q_r g_R^\dagger, \quad g_R \equiv \exp(i Q_R^a r_a). \quad (2.7)$$

The charges are defined by the Noether currents:

$$Q_R^a(t) = \int d^3x J_{R0}^a(x) = \int d^3x q_r^\dagger t^a q_r. \quad (2.8)$$

Using the equal time commutation relations of the field q we obtain the transformation rules of chiral symmetry transformations as

$$\begin{aligned} i [Q_R^a, q_r] &= -it^a q_r, \\ i [Q_L^a, q_l] &= -it^a q_l, \\ i [Q_L^a, q_r] &= i [Q_R^a, q_l] = 0. \end{aligned} \quad (2.9)$$

Furthermore we find the commutation relations among the chiral charges,

$$\begin{aligned} [Q_R^a, Q_R^b] &= i\epsilon_{abc} Q_R^c, \\ [Q_L^a, Q_L^b] &= i\epsilon_{abc} Q_L^c, \\ [Q_R^a, Q_L^b] &= 0. \end{aligned} \quad (2.10)$$

These are the basic commutation relations of the chiral group $SU(2)_R \otimes SU(2)_L$.

It is worth emphasizing that the subscripts R and L are introduced here in order to distinguish the symmetry groups assigned to q_r and q_l ; we could have adopted other labels such as A and B , etc. The important point is that different internal symmetries ($SU(2)$ and $SU(2)$) can be assigned to the right and left handed fermions, independently. This is the reason that we use small letters r and l for the Lorentz group and large letters R and L for the chiral group.

The commutation relations of (2.10) may be equivalently expressed by the vector and axial charge operators,

$$Q_V^a = Q_R^a + Q_L^a, \quad Q_A^a = Q_R^a - Q_L^a, \quad (2.11)$$

which satisfy

$$\begin{aligned} [Q_V^a, Q_V^b] &= i\epsilon_{abc} Q_V^c, \\ [Q_V^a, Q_A^b] &= i\epsilon_{abc} Q_A^c, \end{aligned} \quad (2.12)$$

$$[Q_A^a, Q_A^b] = i\epsilon_{abc} Q_V^c. \quad (2.13)$$

In this VA representation, the right and left handed fermion fields are transformed as

$$\begin{aligned} i [Q_V^a, q_r] &= -it^a q_r, & i [Q_V^a, q_l] &= -it^a q_l, \\ i [Q_A^a, q_r] &= -it^a q_r, & i [Q_A^a, q_l] &= +it^a q_l. \end{aligned} \quad (2.14)$$

In terms of a four component representation, these are combined into

$$i [Q_V^a, q] = -it^a q, \quad i [Q_A^a, q] = -i\gamma_5 t^a q, \quad (2.15)$$

where the γ_5 matrix is diagonal in the chiral (Weyl) representations:

$$\gamma_5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.16)$$

Now we define a notation specifying representations of the chiral group $SU(2)_R \otimes SU(2)_L$ as

$$(I_R, I_L). \quad (2.17)$$

where I_R (I_L) denotes the isospin value of the representation of the group $SU(2)_R$ ($SU(2)_L$). In this notation, the quark field is expressed group theoretically as a direct sum of the fundamental representation of the chiral group:

$$q = q_r + q_l \sim (\frac{1}{2}, 0) \oplus (0, \frac{1}{2}). \quad (2.18)$$

The reason that the quark field is more than an irreducible representation is that it is an eigenstate of a parity transformation.

2.2. Representations for one baryon

After defining chiral symmetry on the quark fields, we construct linear representations for baryons of chiral symmetry from the quark fields. One of the simplest way is to make direct product of quark fields with appropriate quantum numbers for the baryons. For instance, the representation for baryons is given by a direct product of three quarks²⁰⁾

$$\begin{aligned} \psi &\sim q \otimes q \otimes q \sim [(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})]^3 \\ &= 5[(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})] \oplus 3[(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)] \oplus [(\frac{3}{2}, 0) \oplus (0, \frac{3}{2})]. \end{aligned} \quad (2.19)$$

The multiplets $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ and $(\frac{3}{2}, 0) \oplus (0, \frac{3}{2})$ have purely isospin $\frac{1}{2}$ and $\frac{3}{2}$, respectively, while $(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)$ contains both $I = \frac{1}{2}$ and $I = \frac{3}{2}$.

It is natural to choose the $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ representation for the nucleon. This multiplet transforms in the same way as the quark field:

$$i[Q_R^a, \psi_r] = -it^a \psi_r \quad i[Q_L^a, \psi_l] = -it^a \psi_l. \quad (2.20)$$

Here we assume to assign $(\frac{1}{2}, 0)$ to ψ_r . However, it is also possible to assign $(0, \frac{1}{2})$ to ψ_r . We shall discuss this point in the next section.

For Δ , which has $I = \frac{3}{2}$, there are two candidates for the representation of the chiral group (2.19): $[(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)]$ and $[(\frac{3}{2}, 0) \oplus (0, \frac{3}{2})]$. Here we take the former multiplet, since the Δ is known as a strong resonance of $N\pi$ system, and $N \otimes \pi \sim [(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})] \otimes [(\frac{1}{2}, \frac{1}{2})]$ does not contain $(\frac{3}{2}, 0) \oplus (0, \frac{3}{2})$. The transformation rules for a field which belongs to $[(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)]$ are a little bit complicated¹⁶⁾:

$$[Q_R^A, (\psi_r)_i^B] = i\epsilon^{ABC}(\psi_r)_i^C, \quad [Q_R^A, (\psi_l)_i^B] = -(t^A)_{ij}(\psi_l)_j^B, \quad (2.21)$$

$$[Q_L^A, (\psi_r)_i^B] = -(t^A)_{ij}(\psi_r)_j^B, \quad [Q_L^A, (\psi_l)_i^B] = i\epsilon^{ABC}(\psi_l)_i^C. \quad (2.22)$$

The field $(\psi_{r,l})_i^A$ has two indices for the chiral group. Here $A = 1, 2, 3$ is for $I = 1$ triplet, and $i = 1, 2$ for $I = \frac{1}{2}$ doublet. It is important to note that this field contains

both $I = \frac{3}{2}$ and $I = \frac{1}{2}$ components. Therefore Δ and a nucleon resonance N^* form a representation of the chiral group.

From now on in this and the next sections, we do not consider the quark structure of baryons and we specify baryons as linear representations of the chiral group $SU(2)_R \otimes SU(2)_L$. For most part of discussions, we consider the simplest representation of $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ for the nucleon and $(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)$ for the Δ and N^* . In reality, however, other representations could mix with this simplest representation^{4), 5)}. Therefore, results derived in the followings hold to the extent that the mixing is not considered large.

2.3. Representations for two baryons

Let us consider two kinds of baryons, ψ_1 and ψ_2 . Their right and left handed components are denoted as

$$\psi_{1r}, \psi_{1l}, \psi_{2r}, \psi_{2l}. \quad (2.23)$$

To make argument explicit, we consider the nucleon which belongs to $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$. The following argument can be applied to other representations with minor modifications.

We assume that ψ_1 belongs to the same representation as that of (2.9) or (2.15). Namely,

$$i [Q_R^a, \psi_{1r}] = -it^a \psi_{1r}, \quad i [Q_L^a, \psi_{1l}] = -it^a \psi_{1l}. \quad (2.24)$$

Now, for ψ_2 we can assume the same relation

$$i [Q_R^a, \psi_{2r}] = -it^a \psi_{2r}, \quad i [Q_L^a, \psi_{2l}] = -it^a \psi_{2l}. \quad (2.25)$$

We call the set of assignments of (2.24) and (2.25) *naive*^{13), 14)}.

From chiral symmetry alone, however, it is also possible to introduce another assignment for the second fermion,

$$i [Q_L^a, \psi_{2r}] = -it^a \psi_{2r}, \quad i [Q_R^a, \psi_{2l}] = -it^a \psi_{2l}. \quad (2.26)$$

Namely, the right handed fermion transforms under $SU(2)_L$, and the left handed fermion transforms under $SU(2)_R$. A somewhat confusing point stems, as anticipated, from the fact that we have denoted the chiral group as $SU(2)_R \otimes SU(2)_L$ for the right and left handed fermions. Rather, we would have denoted as $SU(2)_A \otimes SU(2)_B$. Since the assignment of the two $SU(2)$ groups is interchanged for the second fermion, we call the assignment (2.24) and (2.26) *mirror*¹⁴⁾.

In the four component representation, we can rewrite the transformation rules of the naive and mirror assignments in the VA representation as ($i = 1, 2$)

$$\text{Naive : } i [Q_V^a, \psi_i] = -it^a \psi_i, \quad i [Q_A^a, \psi_i] = -it^a \gamma_5 \psi_i, \quad (2.27)$$

$$\text{Mirror : } i [Q_V^a, \psi_i] = -it^a \psi_i, \quad i [Q_A^a, \psi_i] = -i\eta t^a \gamma_5 \psi_i, \quad (2.28)$$

where $\eta = 1$ for $i = 1$ and $\eta = -1$ for $i = 2$. These relations imply that the axial charge of ψ_2 in the mirror assignment differs from that of ψ_1 in sign.

In order to distinguish the naive and mirror assignments, we may indicate the chiral group representations of the right and left handed fermions explicitly:

$$\text{Naive : } \psi_1 \sim \psi_{1r}(\tfrac{1}{2}, 0) + \psi_{1l}(0, \tfrac{1}{2}), \quad \psi_2 \sim \psi_{2r}(\tfrac{1}{2}, 0) + \psi_{2l}(0, \tfrac{1}{2}), \quad (2\cdot29)$$

$$\text{Mirror : } \psi_1 \sim \psi_{1r}(\tfrac{1}{2}, 0) + \psi_{1l}(0, \tfrac{1}{2}), \quad \psi_2 \sim \psi_{2r}(0, \tfrac{1}{2}) + \psi_{2l}(\tfrac{1}{2}, 0). \quad (2\cdot30)$$

From Lorentz and chiral invariance, one would have considered combinations

$$\psi_r(\tfrac{1}{2}, 0) + \psi_l(\tfrac{1}{2}, 0) \quad \text{or} \quad \psi_r(0, \tfrac{1}{2}) + \psi_l(0, \tfrac{1}{2}). \quad (2\cdot31)$$

However, this assignment does not allow a Yukawa type meson-baryon interaction term. The chiral field is introduced with the transformation rule as defined in (3\cdot1). If two baryons are defined as the former of (2\cdot31) and are transformed by g_R alone, it is obvious that the interaction term $\bar{\psi}(\sigma + i\vec{\tau}\cdot\vec{\pi}\gamma_5)\psi$ is not chiral invariant. If ψ_1 is defined by the former and ψ_2 by the latter of (2\cdot31), a Lorentz and chiral invariant term seems to be possible:

$$\psi_{2r}^\dagger U \psi_{1l} + \psi_{2l}^\dagger U \psi_{1r} + \psi_{1l}^\dagger U^\dagger \psi_{2r} + \psi_{1r}^\dagger U^\dagger \psi_{2l}, \quad (2\cdot32)$$

where $U = \sigma + i\vec{\tau}\cdot\vec{\pi}$. This term, however, is not invariant under the parity transformation, $\psi_{12,r} \leftrightarrow \psi_{12,l}$ and $U \leftrightarrow U^\dagger$. In this way, we see that the naive and mirror assignments are the two possible chiral assignments for meson and baryon systems.

The transformation rules (2\cdot28) for two isodoublet nucleons were originally introduced in the textbook by Lee already thirty years ago¹⁸⁾. There he constructed a linear sigma model based on that assignment. However his model produced an unphysical consequence $g_{\pi NN} = 0$, since not all relevant terms allowed by chiral symmetry were considered in his Lagrangian. Then DeTar and Kunihiro established a linear sigma model by adding terms missing in the Lee's lagrangian and resolved the problem there. Their model explained a possible parity doubling of massive nucleons¹⁹⁾. The DeTar-Kunihiro model was used in a study of finite density properties of nuclear matter by Hatsuda and Prakash²¹⁾. Beane considered a similar model to the DeTar-Kunihiro model in a different context in ref.²²⁾, where the mirror type assignment is applied to quark fields. Christos also studied parity doubling of nucleon using a local baryon field composed quark fields²³⁾.

2.4. Mass and parity eigenstates

Let us consider a simple lagrangian for one iso-doublet nucleon,

$$L = \sum_{f=p,n} (\bar{\psi}_f i \not{\partial} \psi_f - m \bar{\psi}_f \psi_f), \quad (2\cdot33)$$

where the sum over nucleon isospin is taken. While the kinetic term preserves the chirality, the mass term does not (here we omit writing the flavor indices):

$$\begin{aligned} L_{mass} &= -m \psi_r^\dagger \psi_l - m \psi_l^\dagger \psi_r \\ &\rightarrow -m \psi_r^\dagger g_R^\dagger g_L \psi_l - m \psi_l^\dagger g_L^\dagger g_R \psi_r \quad (\text{not invariant}), \end{aligned}$$

where $g_R \in SU(2)_R$ and $g_L \in SU(2)_R$ are finite group elements of the chiral group transformations as defined in (2\cdot7).

Now we turn to the case of two iso-doublet baryons and consider a simple lagrangian:

$$L = L_K + L_M, \quad (2.34)$$

where the kinetic term is given simply by the sum

$$\bar{\psi}_1 i \not{\partial} \psi_1 + \bar{\psi}_2 i \not{\partial} \psi_2. \quad (2.35)$$

From the Lorentz invariance and baryon number conservation, possible forms of L_M are

$$\begin{aligned} L_M &= -m_1 \left(\psi_{1r}^\dagger \psi_{1l} + \psi_{1l}^\dagger \psi_{1r} \right) - m_2 \left(\psi_{2r}^\dagger \psi_{2l} + \psi_{2l}^\dagger \psi_{2r} \right) \\ &\quad - m_0 \left(\psi_{1r}^\dagger \psi_{2l} + \psi_{2l}^\dagger \psi_{1r} + \psi_{2r}^\dagger \psi_{1l} + \psi_{1l}^\dagger \psi_{2r} \right) \\ &= -m_1 \bar{\psi}_1 \psi_1 - m_2 \bar{\psi}_2 \psi_2 - m_0 (\bar{\psi}_1 \psi_2 + \bar{\psi}_2 \psi_1). \end{aligned} \quad (2.36)$$

For the naive baryons, all the terms break chiral symmetry. Therefore, if we impose the theory to be explicitly chiral invariant, the baryons must be massless.

For the mirror assignment, however, the third term of (2.36) preserves chiral symmetry; for instance,

$$\psi_{1r}^\dagger \psi_{2l} \rightarrow (\psi_{1r}^\dagger g_R^\dagger)(g_R \psi_{2l}) = \psi_{1r}^\dagger \psi_{2l}. \quad (\text{invariant}) \quad (2.37)$$

Because of this, the mirror baryons can acquire a (Dirac) mass keeping chiral symmetry. Since the mass term is off-diagonal in the particle space of ψ_1 and ψ_2 , the physical mass eigenstates are their linear combinations:

$$\psi_+ = \frac{1}{\sqrt{2}}(\psi_1 + \psi_2), \quad \psi_- = \frac{1}{\sqrt{2}}(\psi_1 - \psi_2), \quad (2.38)$$

whose eigenvalues are $\pm m_0$. The eigenvalue of ψ_- appears negative, since ψ_- carries opposite parity of ψ_+ . Multiplying γ_5 , the state $\gamma_5 \psi_-$ carries the same parity as ψ_+ and the eigenvalue turns to $+m_0$ (see Eq. (3.3)).

Now we write the chiral transformation property in terms of $\psi_\pm$:

$$i[Q_V^a, \psi_\pm] = -it^a \psi_\pm, \quad i[Q_A^a, \psi_\pm] = -it^a \gamma_5 \psi_\mp. \quad (2.39)$$

The second equation indicates that the mass eigenstates $\psi_\pm$ form a large representation of the chiral group. It is, however, reducible, since the irreducible representations of the chiral group can not diagonalize the mass matrix. This is a unique feature of the mirror assignment and is a consequence that the eigenstates of chiral symmetry ($\psi_{1,2}$) differs from mass eigenstates, a situation analogous to quark mixing in the standard theory.

§3. Linear Sigma Models

In this section we consider linear sigma models for mesons and nucleons (denoted as N). The meson fields are introduced as components of the representation $(\frac{1}{2}, \frac{1}{2})$ of the chiral group, which are subject to the transformation rule:

$$\sigma + i\vec{\tau} \cdot \vec{\pi} \rightarrow g_L(\sigma + i\vec{\tau} \cdot \vec{\pi})g_R^\dagger. \quad (3.1)$$

The explicit symmetry breaking is not included in the following discussion.

For later convenience, we note the following two properties. First, the parity of the nucleon N changes when multiplied by γ_5 . This is because γ_5 anticommutes with γ_0 :

$$\text{If } P : N \rightarrow \gamma_0 N, \text{ then } P : \gamma_5 N \rightarrow -\gamma_0 \gamma_5 N. \quad (3.2)$$

Second, the Dirac equation for $\gamma_5 N$ changes the sign of m ;

$$(i\partial - m)N = 0 \rightarrow (i\partial + m)\gamma_5 N = 0. \quad (3.3)$$

3.1. The Gell-Mann Levy model

Before going to the linear sigma models with two nucleons, we briefly review some relevant aspects of the Gell-Mann Levy model with one nucleon belonging to $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})^{10)}$. The Lagrangian is given by

$$\mathcal{L}_{GL} = i\bar{N}\not{\partial}N - g\bar{N}(\sigma + i\gamma_5\vec{\tau} \cdot \vec{\pi})N + L_{\text{mes}}, \quad (3.4)$$

where L_{mes} denotes the Lagrangian for σ and π . The explicit form of L_{mes} is irrelevant in the following discussion as long as it causes spontaneous breaking of chiral symmetry. The πNN coupling is given simply by g at the tree level. Once the spontaneous breakdown takes place, the mass of the nucleon is generated as

$$m_N = g\sigma_0, \quad (3.5)$$

where $\sigma_0 \equiv \langle 0|\sigma|0 \rangle$ is the vacuum condensate of the sigma meson. This equation is regarded as the Goldberger-Treiman relation with $g_A = 1$.

In the linear sigma model, it is allowed to add chirally symmetric terms containing derivatives and multi-mesons, once we abandon the renormalizability. For instance, one can add a term like

$$\alpha\bar{N}[(\vec{t} \cdot \vec{\pi}\not{\partial}\sigma - \sigma\vec{t}\not{\partial}\vec{\pi})\gamma_5 - \vec{t} \cdot (\vec{\pi} \otimes \not{\partial}\vec{\pi})]N. \quad (3.6)$$

With this term g_A is modified to be $1 - \alpha\sigma_0^2$ ¹⁸⁾.

Terms with multi-mesons give corrections to the nucleon mass m_N and the pion nucleon coupling g as well as the axial charge in powers of σ_0 . For instance, a chiral-invariant term $\bar{N}(\sigma^2 + \pi^2)^n(\sigma + i\gamma_5\vec{\tau} \cdot \vec{\pi})N$ gives contributions of order σ_0^n and $n\sigma_0^{n-1}$ to m_N and g , respectively. This should be contrasted with the non-linear sigma model which provides contributions in inverse powers of σ_0 . Therefore, in the linear sigma model, the restoration of chiral symmetry is taken smoothly as a function of $\sigma_0 \rightarrow 0$, where correction terms disappear. From the experimental value $g_A^{\text{exp}} = 1.25$, the correction to g_A from the higher derivative terms is of order of several tens percent.

3.2. Naive model

First we consider the naive assignment. It is not difficult to verify that up to forth order in dimension of mass, the general lagrangian can be written as¹⁴⁾:

$$\begin{aligned} L_{\text{naive}} = & \bar{N}_1 i\not{\partial}N_1 - g_1\bar{N}_1(\sigma + i\gamma_5\vec{\tau} \cdot \vec{\pi})N_1 \\ & + \bar{N}_2 i\not{\partial}N_2 - g_2\bar{N}_2(\sigma + i\gamma_5\vec{\tau} \cdot \vec{\pi})N_2 \\ & - g_{12}\{\bar{N}_1(\gamma_5\sigma + i\vec{\tau} \cdot \vec{\pi})N_2 - \bar{N}_2(\gamma_5\sigma + i\vec{\tau} \cdot \vec{\pi})N_1\} + L_{\text{mes}}, \end{aligned} \quad (3.7)$$

where g_1 , g_2 and g_{12} are free parameters. The terms of g_1 and g_2 are ordinary chiral invariant coupling terms of the linear sigma model. The term of g_{12} is the mixing of N_1 and N_2 . Since the two nucleons have opposite parities, γ_5 appears in the coupling with σ , while it does not in the coupling with π .

Chiral symmetry breaks down spontaneously when the sigma meson acquires a finite vacuum expectation value, $\sigma_0 \equiv \langle 0|\sigma|0\rangle$. This generates masses of the nucleons as in (3.5). From (3.7), the mass can be expressed by a $2 \otimes 2$ matrix in the space of N_1 and N_2 . Due to the mixing of different parity states, the diagonalization has to be done carefully. For this we introduce the new fields

$$(N'_1, N'_2) \equiv (N_1, \gamma_5 N_2), \quad (3.8)$$

in terms of which the original lagrangian can be written as:

$$\begin{aligned} L_{\text{naive}} = & \bar{N}'_1 i \not{\partial} N'_1 - g_1 \bar{N}'_1 (\sigma + i \gamma_5 \vec{\tau} \cdot \vec{\pi}) N'_1 \\ & + \bar{N}'_2 i \not{\partial} N'_2 - g_2 \bar{N}'_2 (\sigma + i \gamma_5 \vec{\tau} \cdot \vec{\pi}) N'_2 \\ & - g_{12} \{ \bar{N}'_1 (\sigma + i \gamma_5 \vec{\tau} \cdot \vec{\pi}) N'_2 + \bar{N}'_2 (\sigma + i \gamma_5 \vec{\tau} \cdot \vec{\pi}) N'_1 \} + \dots \end{aligned} \quad (3.9)$$

We note here the sign changes in the g_2 term and the second term of g_{12} terms. The interaction terms, including the mass term, are now expressed in the matrix form as

$$(\bar{N}'_1, \bar{N}'_2) \sigma_0 U_5 \begin{pmatrix} g_1 & g_{12} \\ g_{12} & -g_2 \end{pmatrix} \begin{pmatrix} N'_1 \\ N'_2 \end{pmatrix}, \quad (3.10)$$

where we have introduced $U_5 = (\sigma + i \gamma_5 \vec{\tau} \cdot \vec{\pi})/\sigma_0$, when $\sigma_0 \neq 0$. This term can be diagonalized as

$$(\bar{N}'_+, \bar{N}'_-) \sigma_0 U_5 \begin{pmatrix} g_+ & 0 \\ 0 & -g_- \end{pmatrix} \begin{pmatrix} N'_+ \\ N'_- \end{pmatrix}, \quad (3.11)$$

where

$$g_{\pm} = \frac{1}{2} \left(\sqrt{(g_1 + g_2)^2 + 4g_{12}^2} \pm (g_1 - g_2) \right), \quad (3.12)$$

and

$$\begin{pmatrix} N'_+ \\ N'_- \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} N'_1 \\ N'_2 \end{pmatrix}. \quad (3.13)$$

The mixing angle θ is defined by

$$\tan 2\theta = \frac{2g_{12}}{g_1 + g_2}. \quad (3.14)$$

Turning back to the original fields, $N_{1,2}$, we find the physical eigenfunctions:

$$\begin{pmatrix} N_+ \\ N_- \end{pmatrix} = \begin{pmatrix} \cos \theta & \gamma_5 \sin \theta \\ -\gamma_5 \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} N_1 \\ N_2 \end{pmatrix}, \quad (3.15)$$

whose masses are given by

$$m_{\pm} = \sigma_0 g_{\pm} = \frac{\sigma_0}{2} \left(\sqrt{(g_1 + g_2)^2 + 4g_{12}^2} \pm (g_1 - g_2) \right). \quad (3.16)$$

What is interesting here is that after the diagonalization, the two nucleons decouple completely. Explicitly, the lagrangian can be written as

$$\begin{aligned} L_{\text{naive}} = & \bar{N}_+ i \not{\partial} N_+ - g_+ \bar{N}_+ (\sigma + i\gamma_5 \vec{\tau} \cdot \vec{\pi}) N_+ \\ & + \bar{N}_- i \not{\partial} N_- - g_- \bar{N}_- (\sigma + i\gamma_5 \vec{\tau} \cdot \vec{\pi}) N_- + L_{\text{mes}}. \end{aligned} \quad (3.17)$$

Therefore, chiral symmetry puts no constraint on the relation between N_+ and N_- . The role of chiral symmetry is just the mass generation due to its spontaneous breaking. As shown in Fig. 2 (b), when chiral symmetry is restored and $\sigma_0 \rightarrow 0$, both N_+ and N_- become massless and degenerate. However, the degeneracy is trivial and has no relevance to chiral symmetry between N_+ and N_- . The decoupling of N_+ and N_- implies that the off-diagonal Yukawa coupling $g_{\pi N_+ N_-}$ vanishes in the soft pion limit. This is valid up to the order we considered and if derivative coupling terms are neglected. The same result was also obtained in the QCD sum rule¹²⁾.

3.3. Mirror model

In the mirror assignment of chiral symmetry, the chiral invariant lagrangian up to the order of (mass)⁴ is given by

$$\begin{aligned} L_{\text{mirror}} = & \bar{N}_1 i \not{\partial} N_1 - g_1 \bar{N}_1 (\sigma + i\gamma_5 \vec{\tau} \cdot \vec{\pi}) N_1 \\ & + \bar{N}_2 i \not{\partial} N_2 - g_2 \bar{N}_2 (\sigma - i\gamma_5 \vec{\tau} \cdot \vec{\pi}) N_2 \\ & - m_0 (\bar{N}_1 \gamma_5 N_2 - \bar{N}_2 \gamma_5 N_1) + L_{\text{mes}}. \end{aligned} \quad (3.18)$$

In the g_2 term, the sign of the pion field is opposite to that of the g_1 term, which ensures the chiral invariance of the term in the mirror assignment. This lagrangian was first formulated in the present form by DeTar and Kunihiro^{*)}.

When chiral symmetry is spontaneously broken, the mass matrix of the lagrangian (3.18) is given by

$$M = \begin{pmatrix} g_1 \sigma_0 & m_0 \gamma_5 \\ -m_0 \gamma_5 & g_2 \sigma_0 \end{pmatrix}. \quad (3.19)$$

This can be diagonalized by a linear combination similar to (3.15) with the mixing angle:

$$\tan 2\theta = \frac{2m_0}{\sigma_0(g_1 + g_2)}. \quad (3.20)$$

The corresponding masses of the two states are given by

$$m_{\pm} = \frac{1}{2} \left(\sqrt{(g_1 + g_2)^2 \sigma_0^2 + 4m_0^2} \pm (g_1 - g_2) \sigma_0 \right). \quad (3.21)$$

^{*)} There is a slight difference in the definition in coupling constants.

An interesting feature is that the two nucleons become degenerate with a finite mass m_0 when chiral symmetry is restored.

The mirror model contains richer physics than the naive model, since the diagonalization survives couplings between N_+ and N_- . It is instructive to see first the axial charges which are then related to the one pion Yukawa couplings through the Goldberger-Treiman relation. The axial charges are extracted from the commutation relations:

$$[Q_A^a, N_+] = \frac{\tau^a}{2} \gamma_5 (\cos 2\theta N_+ - \sin 2\theta \gamma_5 N_-), \quad (3.22)$$

$$[Q_A^a, N_-] = \frac{\tau^a}{2} \gamma_5 (-\sin 2\theta \gamma_5 N_+ - \cos 2\theta N_-), \quad (3.23)$$

giving a matrix expression

$$g_A = \begin{pmatrix} \cos 2\theta & -\sin 2\theta \gamma_5 \\ -\sin 2\theta \gamma_5 & -\cos 2\theta \end{pmatrix}. \quad (3.24)$$

We would like to make several comments.

1. As anticipated, the axial charge of the negative parity nucleon is just the minus of that of the positive parity nucleon. This feature holds at any mixing angle θ .
2. The absolute value of the diagonal component is, however, less than unity. This may be increased by including higher derivative terms.
3. The present result obtained from the linear sigma model can be generalized by a group theoretical method, where the nucleon axial charge $g_A \sim \langle N | Q_A^a | N \rangle$ is computed from the commutation relation $[Q_A^+, Q_A^-] = -Q_V^{(4), 5)}$. Assuming that the intermediate states of

$$\langle N | Q_A^+ Q_A^- | N \rangle = \sum_B \langle N | Q_A^+ | B \rangle \langle B | Q_A^- | N \rangle. \quad (3.25)$$

are saturated by one baryon states, the method reproduces the result obtained in the linear sigma model.

Now we extract the one pion Yukawa couplings from the lagrangian written in terms of the physical basis N_+ and N_- . The results are

$$g_{\pi N_+ N_+} = g_1 \cos^2 \theta + g_2 \sin^2 \theta = \frac{m_+}{\sigma_0} g_A^{++}, \quad (3.26)$$

$$g_{\pi N_- N_-} = -(g_1 \sin^2 \theta + g_2 \cos^2 \theta) = \frac{m_-}{\sigma_0} g_A^{--}, \quad (3.27)$$

$$g_{\pi N_+ N_-} = \frac{g_2 - g_1}{2} \sin 2\theta = \frac{m_- - m_+}{2\sigma_0} g_A^{+-}. \quad (3.28)$$

The second equalities of these equations are the generalized Goldberger-Treiman relations.

Finally we perform a simple parameter fitting¹⁹⁾. In doing so, we assume to identify $N_+ \rightarrow N(939) \equiv N$ and $N_- \rightarrow N(1535) \equiv N^*$. In order to fix the four parameters, we use the inputs $m_+ = 939$, $m_- = 1535$ MeV, $\sigma_0 = f_\pi = 93$ MeV

and $g_{\pi N_+ N_-} \sim 0.7$. The last equation is deduced from the partial decay width of $N(1535)^{11)}$ (although large uncertainties for the width have been reported^{33), 34)}:

$$\Gamma_{N^* \rightarrow \pi N} \sim 75 \text{ MeV}, \quad (3.29)$$

and the formula

$$\Gamma_{N^* \rightarrow \pi N} = \frac{3q(E_N + M)}{4\pi M^*} g_{\pi N N^*}^2, \quad (3.30)$$

where q is the relative momentum in the final state. We find

$$\begin{aligned} \sigma_0 &= 93 \text{ MeV}, \quad m_0 = 270 \text{ MeV}, \\ g_1 &= 9.8, \quad g_2 = 16. \end{aligned} \quad (3.31)$$

From these parameters, we find the mixing angle

$$\theta = 6.3 \text{ degree}, \quad (3.32)$$

giving the diagonal value of the axial charges

$$g_A^{++} = -g_A^{--} = 0.98. \quad (3.33)$$

The different sign of the g_A 's as indicated in (3.33) leads to different contributions in the calculation of the anomalous triangle diagram, when the loops of N and N^* are considered. The result may contradict if the anomaly matching is required as an effective theory of QCD. However, it is argued that the anomaly should be accounted by quark loops rather than baryon loops and that the effective lagrangian emerges when quark loops are integrated out. Therefore, although not shown in our present lagrangians, the Wess-Zumino-Witten term should be added in order to account for the correct anomaly. In the mirror model, this interpretation is rather welcome, since baryon loop contributions of N and N^* cancel due to the opposite signs of the g_A 's and the entire anomaly is attributed to the quark loops. In this regard, the naive model is rather in contradiction as far as the anomaly matching to QCD is concerned, since baryon loops contribute constructively with the same sign of the g_A 's.

3.4. Toward chiral symmetry restoration

The change in chiral symmetry is dictated by the order parameter σ_0 , which is identified with the pion decay constant f_π . It decreases as temperature or density is increased. The naive and mirror models exhibit different behaviors of physical quantities such as masses and coupling constants as chiral symmetry is restored. The behavior of the naive case is rather trivial, and therefore, here we discuss mostly the mirror case.

- Mixing angles and masses (Figs. 2)

When chiral symmetry is restored ($\sigma_0 = 0$), the only source of the mixing is the off-diagonal mass term (see (2.36) or (3.19)). Therefore, the two degenerate states ψ_1 and ψ_2 mix with the equal weight ($\theta = \pi/4$) having an equal mass




Fig. 2. (a) Mixing angle of the mirror model, and (b) the masses of the positive and negative parity nucleons in the naive and mirror models, as functions of the chiral condensate σ_0 .




Fig. 3. Diagonal and off-diagonal axial charges in the naive and mirror models, as functions of the chiral condensate σ_0 .

$m_{\pm} = m_0$. As the interaction is turned on and chiral symmetry starts to break spontaneously, the mixing angle decreases monotonically and reaches 6.3 degree in the real world ($\sigma_0 = 93$ MeV). Also, as f_{π} is increased the masses increase and the degeneracy is resolved: $m_+ = 939$ MeV and $m_- = 1535$ MeV when $\sigma_0 = 93$ MeV.

- g_A (Figs. 3)

It is interesting to see the behavior of the diagonal $g_A^{++} = -g_A^{--}$ and off-diagonal g_A^{+-} as functions of σ_0 . In particular, g_A^{+-} increases as chiral symmetry starts to restore. Since the axial charges are related to pion couplings, the increase of the off-diagonal couplings $g_A^{+-} \sim g_{\pi N_+ N_-}$ was considered as a cause of the increase in the width of $N(1535)$ in medium^{17), 27), 28)}. More detailed calculation was reported in Ref.¹⁷⁾.

3.5. Higher representations

Recently, Jido, Hatsuda and Kunihiro extended the simple mirror model of fundamental representations to a model of higher dimensional representations¹⁶⁾. They considered a representation of $(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)$. The physics motivation of adopting it lies in the fact that the pion couples strongly with the nucleon and delta. A simple

irreducible decomposition

$$\left(\left(\frac{1}{2}, 0\right) \oplus \left(0, \frac{1}{2}\right)\right) \otimes \left(\frac{1}{2}, \frac{1}{2}\right) = \left(\left(1, \frac{1}{2}\right) \oplus \left(\frac{1}{2}, 1\right)\right) \oplus \left(\left(0, \frac{1}{2}\right) \oplus \left(\frac{1}{2}, 0\right)\right). \quad (3.34)$$

implies that the higher dimensional representation is realized by a composite system of the pion and nucleon. Incidentally, it is interesting to recognize that the second term of (3.34) is the mirror representation of the original nucleon. This is the basis of the five quark representation for the mirror baryon as discussed in the next section.

Since the representation $(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)$ contains both isospin 1/2 and 3/2 components, they have reached a *quartet scheme* in which Δ , N^* and their parity partners form the chiral multiplet. Although their argument can be applied for baryons with any spin J , to make argument explicit, here we take baryons with spin $J = \frac{3}{2}$ as an example. Then in the quartet scheme we have $\Delta_+(P_{33})$, $\Delta_-(D_{33})$, $N_+^*(P_{13})$ and $N_-^*(D_{13})$.

In the mirror assignment, ψ_{1r} and ψ_{2l} belong to $(1, \frac{1}{2})$, while ψ_{1l} and ψ_{2r} belong to $(\frac{1}{2}, 1)$. Here the Lorentz index for the Rarita-Schwinger field with $J = \frac{3}{2}$ is omitted for simplicity. The transformation rules under the chiral rotations are given by

$$\begin{aligned} [Q_R^A, (\psi_{1r,2l})_i^B] &= i\epsilon^{ABC}(\psi_{1r,2l})_i^C, \\ [Q_R^A, (\psi_{1l,2r})_i^B] &= -(t^A)_{ij}(\psi_{1l,2r})_j^B, \\ [Q_L^A, (\psi_{1r,2l})_i^B] &= -(t^A)_{ij}(\psi_{1r,2l})_j^B, \\ [Q_L^A, (\psi_{1l,2r})_i^B] &= i\epsilon^{ABC}(\psi_{1l,2r})_i^C. \end{aligned} \quad (3.35)$$

The reason that the naive assignment was not taken is that it gives an unrealistic mass relation $m_{\Delta\pm} = 2m_{N_\mp^*}$. The way to construct a chiral invariant Lagrangian is similar to what we have explored here¹⁶⁾. The interaction Lagrangian is given by

$$\begin{aligned} \mathcal{L}_{\text{int}} &= m_0 \left(\bar{\psi}_2^A \gamma_5 \psi^A - \bar{\psi}_1^A \gamma_5 \psi_2^A \right) \\ &+ a \bar{\psi}_1^A \tau^B (\sigma - i\gamma_5 \vec{\pi} \cdot \vec{\tau}) \tau^A \psi_1^B + b \bar{\psi}_2^A \tau^B (\sigma + i\gamma_5 \vec{\pi} \cdot \vec{\tau}) \tau^A \psi_2^B. \end{aligned} \quad (3.36)$$

The fields with the particle basis $(\Delta_{1,2}, N_{1,2}^*)$ are obtained by the following isospin decomposition:

$$(\psi_{1,2})_i^A = \sum_M (T_{\frac{3}{2}}^A)_{iM} \Delta_{1,2}^M + \sum_m (T_{\frac{1}{2}}^A)_{im} N_{1,2}^{*m}, \quad (3.37)$$

where the projection matrices are defined through the Clebsch-Gordan coefficients:

$$(T_I^A)_{im} = \sum_{\mu\nu} (1\mu \frac{1}{2}\nu | Im) \epsilon_\mu^A \chi_\nu^i, \quad (3.38)$$

where $(\epsilon_\mu^A) = \vec{\epsilon}_\mu$ are polarization vectors

$$\vec{\epsilon}_{+1} = -\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \\ 0 \end{pmatrix}, \quad \vec{\epsilon}_{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \\ 0 \end{pmatrix}, \quad \vec{\epsilon}_0 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (3.39)$$

Table I. Comparison between parameter free predictions of the quartet scheme (QS) and the data¹⁶⁾. Case 1 and case 2 in the $J = \frac{1}{2}$ sector stand for the cases where $N_-^* = N(1535)$ and $N_-^* = N(1650)$, respectively. The last two rows are model parameters m_0, a and b determined from the experimental inputs.

	QS	$J = \frac{1}{2}$		$J = \frac{3}{2}$	$J = \frac{5}{2}$
		case 1	case 2		
$\text{sgn}\left(\frac{m_{N_+^*} - m_{N_-^*}}{m_{\Delta_+} - m_{\Delta_-}}\right)$	—	—	—	—	—
$\frac{m_{N_+^*} - m_{N_-^*}}{m_{\Delta_+} - m_{\Delta_-}}$	$-\frac{1}{2}$	-0.33	-0.72	-0.43	-0.2
$\frac{m_{N_+^*} + m_{N_-^*}}{m_{\Delta_+} + m_{\Delta_-}}$	≤ 1	0.84	0.88	1.1	0.87
m_0 (MeV)		1380	1460	1540	1590
(a, b)		(5.2, 6.6)	(4.4, 6.1)	(1.2, -1.2)	(5.8, 5.7)

and $(\chi_\nu^i) = \chi_\nu$ spinors

$$\chi_{1/2} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \chi_{-1/2} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (3.40)$$

After spontaneous breakdown of chiral symmetry and diagonalization of the mass matrices, the masses of the Δ 's and N^* 's are given by

$$m_{\Delta_\pm} = \sqrt{(a+b)^2 \sigma_0^2 + m_0^2} \mp \sigma_0(a-b), \quad (3.41)$$

$$m_{N_\pm^*} = \sqrt{\left(\frac{a+b}{2}\right)^2 \sigma_0^2 + m_0^2} \pm \frac{\sigma_0}{2}(a-b). \quad (3.42)$$

These equations show that the chiral symmetry breaking induces the mass splitting between parity partners as well as between isospin states. It is interesting to note that these equations lead to parameter free and nontrivial relations among baryon masses:

$$\begin{aligned} \text{sgn}(m_{\Delta_+} - m_{\Delta_-}) &= -\text{sgn}(m_{N_+^*} - m_{N_-^*}), \\ \frac{1}{2}(m_{\Delta_-} - m_{\Delta_+}) &= m_{N_+^*} - m_{N_-^*}, \\ \frac{1}{2}(m_{\Delta_+} + m_{\Delta_-}) &\geq \frac{1}{2}(m_{N_+^*} + m_{N_-^*}). \end{aligned} \quad (3.43)$$

As mentioned before, the present argument holds for the resonances with an arbitrary spin as far as we consider the $(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)$ multiplet. Here we assign the quartet to the lowest resonances in each spin parity among the observed baryons. However, for $J = \frac{1}{2}$ the nucleon is assumed to belong to $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ with a possible parity partner, $N(1535)$ or $N(1650)$ (or even their linear combinations). Therefore, we investigate two quartet schemes for $J = \frac{1}{2}$: that including $N(1535)$ (case 1) or that including $N(1650)$ (case 2). In table I, the parameter free relations (3.43) are compared with those expected from observed masses.

Encouraged by the agreement of the parameter free relations, we determine three model parameters. This is done for each quartet of different spin. We take




Fig. 4. The quartet members with $J = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}$ (¹⁶). The right (left) hand side for each spin is the observed (quartet scheme) masses. The solid (dashed) lines denote the even (odd) parity baryons. The reproduced masses in our scheme agree with the experimental values within 10 %.

$f_\pi = 93$ MeV, and then four experimental masses are used for least square fitting. The resulting parameters are presented in table I. In Fig.4 masses of the quartet members are compared with experiments¹⁶⁾. It is shown that the experimental masses are fitted within 10%.

In the $J = \frac{3}{2}$ the fitted parameters $a = -b = 1.2$ lead approximately to interesting selection rules for the π -decay of the the quartet baryons. If $a = -b$, the interaction Lagrangian of π and $\psi_\pm$ is written as

$$\mathcal{L}_{1\pi} = (\bar{\psi}_+^A, \bar{\psi}_-^A) \begin{pmatrix} 0 & -a \\ a & 0 \end{pmatrix} \tau^B (i\vec{\pi} \cdot \vec{\tau}) \tau^A \begin{pmatrix} \psi_+^B \\ \psi_-^B \end{pmatrix} \quad (3.44)$$

where $\psi_+ = \frac{1}{\sqrt{2}}(\psi_1 + \gamma_5 \psi_2)$ and $\psi_- = \frac{1}{\sqrt{2}}(\gamma_5 \psi_1 - \psi_2)$. This interaction Lagrangian implies that the parity nonchanging couplings, such as $\pi \Delta_\pm \Delta_\pm$, $\pi \Delta_\pm N_\pm^*$ and $\pi N_\pm^* N_\pm^*$, are forbidden at the tree level. This is qualitatively consistent with the decay patterns of the observed resonances. These suppressions, in fact, have been seen also in the constituent quark model²⁵⁾ and the collective string model²⁶⁾, although the considered symmetry and dynamics are quite different from the present model. Therefore it is interesting to compare the physical reason and the relations among the models to understand the role of chiral symmetry in the baryon resonances.

§4. QCD Spectral Function of Positive and Negative Parity Baryons

We have seen that the low energy effective theory of hadrons is strongly constrained by chiral symmetry assignment of hadrons. As QCD is the origin of the effective theory, the assignment must be consistent with QCD. We face two crucial questions: (1) which assignment is consistent with QCD for chiral symmetry of baryons? and (2) how can we confirm the assignment for actual baryon resonances?

In order to answer these questions, we consider the following two steps. First we

construct local operators (interpolating fields) for hadrons in terms of quark fields as components of a suitable chiral group representation. Then, as the second step, we consider chiral properties of hadron states which are coupled to the local operators. Since chiral symmetry is spontaneously broken and the QCD vacuum is not chiral invariant, the second step is nontrivial. We consider two-point correlation functions of baryons and how the chiral symmetry of the local operators is connected to that of the baryonic states.

4.1. Interpolating fields

Under the $SU(2)_R \otimes SU(2)_L$ chiral transform, the quark operator $q(x)$ behaves as the fundamental representation, $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ (see Eq. (2.15)):

$$[Q_A^a, q(x)] = -\gamma_5 t^a q(x), \quad (4.1)$$

where $t^a = \tau^a/2$ are the generators of $SU(2)$. The quark field is an $SU(2)$ isospin doublet and is denoted as

$$q(x) = \begin{pmatrix} u(x) \\ d(x) \end{pmatrix}.$$

Here the Dirac indices of quark fields and gamma matrices are suppressed.

To start with, let us see how interpolating fields for mesons are constructed as desired chiral group representations. The pion is special as the Nambu-Goldstone boson of spontaneous breaking of chiral symmetry. The PCAC (in the strong sense) conjectures a widely accepted relation,

$$\partial_\mu A^{a\mu}(x) = m_\pi^2 f_\pi \phi_\pi^a(x).$$

Then the pion corresponds to a local quark bilinear operator,

$$\phi_\pi^a(x) \sim \partial_\mu [\bar{q}(x) \gamma^\mu \gamma_5 \tau^a q(x)]_{\text{CS}} \sim 2m_q [\bar{q}(x) i \gamma_5 \tau^a q(x)]_{\text{CS}}.$$

Here $[\bar{q} \dots q]_{\text{CS}}$ means that $(\bar{q} \dots q)$ is made color-singlet by summing over the color indices. It is easy to see that this operator is transformed as the a -th component of a chiral $O(4)$ vector. The axial transformation gives

$$[Q_A^a, [\bar{q} i \gamma_5 \tau^b q]_{\text{CS}}] = i \delta^{ab} [\bar{q} q]_{\text{CS}} \quad : \quad [Q_A^a, [\bar{q} q]_{\text{CS}}] = -i [\bar{q} i \gamma_5 \tau^a q]_{\text{CS}} \quad (4.2)$$

Here $[\bar{q} q]_{\text{CS}}$ is the fourth component of the $O(4)$ vector, which corresponds to a scalar isoscalar meson, i.e. σ . Thus the set of operators,

$$([\bar{q}(x) i \gamma_5 \tau^a q(x)]_{\text{CS}}, [\bar{q}(x) q(x)]_{\text{CS}}) \sim (\pi^a, \sigma), \quad (4.3)$$

belong to a linear representation $(\frac{1}{2}, \frac{1}{2})$ of the chiral $SU(2)_R \otimes SU(2)_L$ group. The mass splitting of π and σ is therefore a manifestation of the spontaneous breaking of chiral symmetry.

Similarly, one can introduce vector mesons as two independent representations: $(1, 0)$ and $(0, 1)$. They are represented by $[\bar{q}_l(x) \gamma^\mu \tau^a q_l(x)]_{\text{CS}}$ and $[\bar{q}_r(x) \gamma^\mu \tau^a q_r(x)]_{\text{CS}}$, where the right and left projections are defined by (2.4). Each of these representation is not an eigenstate of the parity, but the $r \pm l$ combinations give the vector and

axial-vector operators of parity eigenstates. Again in the chiral limit, they must be degenerate, while the physical mesons corresponding to these currents are ρ and a_1 , which are split by about 400 MeV.

Now we consider the nucleon operator, which is supposed to consist of three quark operators. A choice of such an operator is³⁰⁾

$$B^\alpha(x) \equiv \left[(q^T(x) C \gamma_5 q(x))_{I=0} q(x)^\alpha \right]_{\text{CS}} \quad (4.4)$$

where α is the Dirac index of the nucleon and C is the charge conjugation gamma matrix, $C = i\gamma^2\gamma^0$ (standard representation). This is a local operator with the appropriate quantum number for the nucleon, $S = \frac{1}{2}$ and $I = \frac{1}{2}$. The first two quarks in (4.4) form a scalar di-quark with $I = 0$, which commutes with Q_A^a . Thus the nucleon operator itself behaves just like a quark field under chiral transform,

$$[Q_A^a, B(x)] = -\gamma_5 t^a B(x) \quad (4.5)$$

This equation indicates that B belongs to the linear representation, $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$, with the axial charge $g_A = 1$.

There is an alternative choice for the nucleon operator,

$$\tilde{B}^\alpha(x) \equiv \left[(q^T(x) C q(x))_{I=0} \{ \gamma_5 q(x) \}^\alpha \right]_{\text{CS}}. \quad (4.6)$$

It is easy to see that $\tilde{B}$ also satisfies eq.(4.5), and therefore has $g_A = 1$. In the following arguments we need not distinguish $\tilde{B}$ from B . Indeed, most general local operator (without derivative) for the proton is given explicitly by³¹⁾

$$B_p(x) = \epsilon^{abc} \left[(u^a{}^T C d^b) \gamma_5 u^c + t (u^a{}^T C \gamma_5 d^b) u^c \right], \quad (4.7)$$

where a , b , and c stand for the color of the quarks, and t is a parameter which controls the mixing of the two independent operators.

In the previous sections, we have pointed out the possibility that an excited nucleon is a mirror partner of the ground state. The signature of the mirror assignment is the negative axial charge g_A . We have seen that the three-quark baryon operators, (4.4) and (4.6), give the naive nucleon. It turns out that the mirror baryon consists of more than three quarks if it is constructed as a local operator without derivatives.

We consider the color-singlet scalar/pseudoscalar bilinear operators, given by (4.3), which are transformed as (4.2). Then, the mirror baryon can be represented by a local 5-quark operator,

$$B^{*\alpha} = [\bar{q}q]_{\text{CS}} B^\alpha + [\bar{q}i\gamma_5\tau^b q]_{\text{CS}} (i\gamma_5\tau^b B)^\alpha, \quad (4.8)$$

where B is defined by either (4.4) or (4.6). Using eqs.(4.5) and (4.2), one can check

$$[Q_A^a, B^*] = +\gamma_5 t^a B^*, \quad (4.9)$$

and thus B^* has a negative axial charge $g_A = -1$. We conclude that B^* represents a mirror nucleon.

4.2. Correlation functions

If chiral symmetry were realized in the Wigner mode, i.e. not spontaneously broken, in QCD, then the operators with different chirality would not mix and would generate distinct hadron states. Unfortunately (or otherwise) chiral symmetry is broken.

For example, consider matrix elements,

$$\langle 0 | [\bar{q} i \gamma_5 \tau^b q]_{\text{CS}} | \pi^a \rangle, \quad \text{and} \quad \langle 0 | [\bar{q} q]_{\text{CS}} | \sigma \rangle. \quad (4.10)$$

Using eq.(4.2), we find

$$\langle 0 | [\bar{q} i \gamma_5 \tau^a q]_{\text{CS}} | \pi^a \rangle = \langle 0 | i [Q_A^a, [\bar{q} q]_{\text{CS}}] | \pi^a \rangle.$$

In the Wigner mode, $Q_A^a | 0 \rangle = 0$ and (π^a, σ) form a $O(4)$ vector. Therefore, we find

$$\langle 0 | [\bar{q} i \gamma_5 \tau^a q]_{\text{CS}} | \pi^a \rangle = -\langle 0 | i [\bar{q} q]_{\text{CS}} Q_A^a | \pi^a \rangle = +\langle 0 | [\bar{q} q]_{\text{CS}} | \sigma \rangle.$$

Thus, the matrix elements, (4.10), are related to each other. This is, however, not the case if the symmetry is spontaneously broken. Q_5^a does not annihilate the vacuum and the pion and sigma states are not simply related by the axial transformation. Instead, if we apply the soft pion theorem³²⁾, we obtain in the limit $p_\pi \rightarrow 0$,

$$\langle 0 | [\bar{q} i \gamma_5 \tau^a q]_{\text{CS}} | \pi^a \rangle = -\frac{i}{f_\pi} \langle 0 | [Q_A^a, [\bar{q} i \gamma_5 \tau^a q]_{\text{CS}}] | 0 \rangle = \frac{1}{f_\pi} \langle 0 | [\bar{q} q]_{\text{CS}} | 0 \rangle.$$

Thus the pion matrix element is related to the $\bar{q}q$ condensate in the vacuum. This relation leads to the Gell-Mann-Oakes-Renner relation.

In the baryon sector, we expect that the symmetry breaking in the vacuum mixes different chiral representations. Furthermore, it should be noted that the local operator (4.4) or (4.6) does not create only the ground state nucleon but also couples to excited states. It is therefore necessary to single out information of the ground state in order to study the properties of the nucleon. A standard technique to achieve this separation is to consider a two-point correlator of the above operators²⁹⁾,

$$T(p) \equiv i \int d^4x e^{ipx} \theta(x^0) \langle 0 | B(x) \bar{B}(0) | 0 \rangle = \gamma^0 A(\sqrt{s}) + B(\sqrt{s}), \quad (4.11)$$

where we take the reference frame $p^0 = \sqrt{s}$, and $\vec{p} = 0$. It is easy to see that the spectral representation of this correlation function contains contributions of both the positive and negative parity baryons. That is,

$$T(p) = \frac{1}{\pi} \int \frac{1 + \gamma^0}{\sqrt{s} - m - i\epsilon} \rho^{(+)}(m) dm - \frac{1}{\pi} \int \frac{1 - \gamma^0}{\sqrt{s} - m - i\epsilon} \rho^{(-)}(m) dm \quad (4.12)$$

where the first (second) term represents the contribution of the positive (negative) parity baryon spectrum. Using the dispersion relation, we find

$$\rho^{(\pm)}(m) = \frac{1}{2} \text{Im}[A(m) \pm B(m)] \quad (4.13)$$

We immediately see that $\rho^{(\pm)}(m)$ are equal if $B(m) = 0$, while the observed baryon spectrum shows that they are not.

Eq.(4.11) is often employed in constructing QCD sum rules²⁹⁾. The correlation function can be evaluated in the deep Euclid momentum region, i.e., $-(p^0)^2 \rightarrow \infty$, resulting an expression in terms of vacuum expectation values of various local operators, such as the quark condensate and gluon condensate. Then using the dispersion relation, we derive a relation of the spectral function ρ to the QCD vacuum properties.

Applying the QCD sum rule to (4.11) we find that chiral symmetry breaking is responsible for the splitting of positive and negative parity baryons. It turns out that the QCD part of $\text{Im } B(m)$ contains the local operators which break chiral symmetry, such as $\langle \bar{q}q \rangle$, $\langle \bar{q}\sigma_{\mu\nu}G^{\mu\nu}q \rangle$. Thus one sees directly that chiral symmetry breaking makes $B(m)$ nonzero and therefore produces the mass difference of the positive and negative parity baryons.

A careful QCD Borel sum rule analysis with standard values of the quark condensate and other QCD parameters confirms that the spectrum of the positive and negative parity octet baryons,²⁹⁾ i.e., the nucleon, Λ , Σ and Ξ , is fairly well reproduced. It is also shown that the flavor singlet baryon shows a special behavior so that the negative parity state has a lower mass than its positive parity partner. The lower state is assigned to the lightest observed negative-parity strange-baryon $\Lambda(1405)$.

This result shows that the mass splittings of the positive and negative parity baryons are attributed to chiral symmetry breaking. Can we tell whether they are destined to be mirror baryons in chiral symmetry restoration? In Eq.(4.11), we employ the naive baryon operators, i.e. three-quark operators of (4.7). We have seen that their correlation function gives excited baryon spectrum that agrees with experiment reasonably. However, as the chiral symmetry breaking mixes chiral representations in general, we do not immediately conclude that the excited baryons which are coupled to (4.7) are “naive”. Further, we may have more than one baryon resonances in a relatively small energy region, as is for $1/2^- N^*$. This unfortunate situation prevents us from making a definite conclusion on the chiral properties of the observed baryon resonances.

However, we can study the properties of the baryon operators by using the correlation functions. Two point correlation functions of the mirror nucleon operator reveal interesting features on the pion-nucleon couplings. Especially, the off-diagonal coupling of the pion to the positive parity and negative parity nucleons can be shown to vanish for the naive baryons. It can be seen in the soft-pion limit of the two-point correlator with a background pion:

$$\begin{aligned}
T_\pi^a(p) &= \lim_{q \rightarrow 0} i \int d^4x e^{ip \cdot x} \langle 0 | T(B(x)B^*(0)) | \pi^a(q) \rangle \\
&= i(A_\pi^a(p^2)p \cdot \gamma \gamma_5 + B_\pi^a(p^2)\gamma_5) \\
&= -\frac{i}{f_\pi} \int d^4x e^{ip \cdot x} \langle 0 | [Q_5^a, T(B(x)B^*(0))] | 0 \rangle. \tag{4.14}
\end{aligned}$$

The same correlator can be expressed in terms of the πBB^* coupling constant by

$$T_\pi^a(p) \simeq g_{\pi BB^*} \lambda_B \lambda_{B^*} \frac{m_B + m_{B^*}}{(p^2 - m_B^2)(p^2 - m_{B^*}^2)} i p \cdot \gamma \gamma_5 \tau^a + \dots \quad (4.15)$$

for the negative parity baryon excitation B^* . This shows that $A(p^2)$ in (4.14) is the term responsible for the off-diagonal πBB^* coupling. Substituting the 3-quark or 5-quark operators for B and B^* in (4.14), we find that for the naive B^* the term A vanishes identically in the soft-pion limit¹²⁾, while for the mirror B^* it is related to the chiral-even term of the BB^* correlator in the vacuum.

This result is consistent with the linear sigma model described in section 3. Namely, in the naive case, as the π -baryon coupling comes from the term that is responsible for the baryon masses, the coupling is diagonal for the mass eigenstate of baryons.

In conclusion, further studies of the baryonic correlation functions, both in QCD sum rules and in lattice QCD calculations will be useful. It is, however, not straightforward to answer the questions like whether a particular nucleon excited state is a mirror of the ground state (or the other excited state) baryon and how much different chiral symmetry representations are mixed in the excited states.

§5. Observation of Chiral Properties of Baryons

5.1. Introduction

In this section, we study possible reactions to distinguish our two models for nucleons, which have been discussed in the previous sections. For this we propose two meson productions of π and η at the threshold region induced by either the pion or photon.

As discussed in the preceding sections, one of the differences between the naive and mirror assignments is the relative sign of the axial charges of the positive and negative parity nucleons. In the following discussions, we identify $N_+ \sim N(939) \equiv N$ and $N_- \sim N(1535) \equiv N^*$. This identification for the nucleons is no more than an assumption. From experimental point of view, however, $N(1535)$ has an advantage that it has a strong coupling to an η meson, which can be used as a signature for the resonance.

In the reactions we propose, the relative sign of the axial charges may be observed by interference effects. In fact, the axial charges are related to the pion Yukawa couplings through the Goldberger-Treiman relations (3.26) and (3.27). Hence the relative sign of $g_{\pi NN}$ and $g_{\pi N^* N^*}$ follows that of $g_A^{NN} (\equiv g_A^{++})$ and $g_A^{N^* N^*} (\equiv g_A^{--})$. The essential idea may be explained by using two diagrams shown in Fig. 5. The incident particle can be either the pion or photon. Let us assume that the two diagrams dominate the reaction close to the threshold. The diagrams (1) and (2) in Fig. 5 have the coupling $g_{\pi NN}$ and $g_{\pi N^* N^*}$, respectively. Therefore the two terms interfere either constructively or destructively, depending on the relative sign of the couplings. In reality, due to kinematical conditions, other diagrams will contribute, but we will find that there are some cross sections in which interference effect can be seen clearly.




Fig. 5. Two diagrams for π and η productions. The incident particle denoted by wavy line is either pion or photon.




Fig. 6. Six N^* dominant diagrams for $\pi^- p \rightarrow \pi^- \eta p$ process.

5.2. Pion induced reaction: $\pi + N \rightarrow \pi + \eta + N$

First we investigate the pion induced process. Here we consider $p\pi^- \rightarrow p\pi^-\eta$ process, since there are two charged particles in the final state, which has an advantage in the experimental setup. If the emitted proton and pion are observed and their invariant mass is measured, it is not necessary to observe the neutral eta meson. $p\pi^- \rightarrow n\pi^0\eta$ is also possible, in principle, to observe the relative sign, but there are more neutral particles. $p\pi^+ \rightarrow p\pi^+\eta$ cannot be used, since $p\pi^+$ state has purely isospin $I = 3/2$ and cannot couple with $N(1535)$ before emitting a pion. Therefore there is no chance to enter $g_{\pi N^* N^*}$ in the process.

We assume $N^* = N(1535)$ dominance as in single η productions, where it is known that the η couples strongly between N and N^* , while other couplings are negligibly small. In this N^* dominance, we can write altogether six diagrams as shown in Figs. 6.

If we strictly follow the linear sigma model lagrangians in section 3, the diagrams involving $\pi N N^*$ vanishes for the naive assignment. However, if we consider small explicit breaking of chiral symmetry and/or higher order derivative terms, corrections yield the small coupling $g_{\pi N N^*} \sim 0.7$ which is extracted from the decay width $\Gamma_{N^* \rightarrow \pi N} \sim 75$ MeV. is introduced in the naive. Hence in this section we consider the difference of the naive and the mirror model only in the relative sign of the $\pi N N$ and $\pi N^* N^*$ coupling constants.

In this spirit, we adopt the following interaction lagrangians for actual computations of cross sections:

$$L_{\pi NN} = g_{\pi NN} \bar{N} i \gamma_5 \vec{\tau} \cdot \vec{\pi} N,$$

Table II. Parameters used in our calculation.

m_N	m_{N^*}	Γ_{N^*}	$g_{\pi NN}$	$g_{\pi NN^*}$	$g_{\eta NN^*}$	$g_{\pi N^* N^*}$
938	1535	140	13	0.7	2.0	13 (naïve)
(MeV)	(MeV)	(MeV)				-13 (mirror)

$$\begin{aligned}
L_{\eta NN^*} &= g_{\eta NN^*}(\bar{N}\eta N^* + \bar{N}^*\eta N), \\
L_{\pi NN^*} &= g_{\pi NN^*}(\bar{N}\tau \cdot \pi N^* + \bar{N}^*\tau \cdot \pi N), \\
L_{\pi N^* N^*} &= g_{\pi N^* N^*}(\bar{N}^*i\gamma_5\tau \cdot \pi N^*).
\end{aligned} \tag{5.1}$$

We use empirical values for coupling constant; $g_{\pi NN} \sim 13$, $g_{\pi NN^*} \sim 0.7$ and $g_{\eta NN^*} \sim 2$. The unknown parameter is the $g_{\pi N^* N^*}$ coupling. One can estimate it by using the theoretical value of the axial charge $g_A^{N^* N^*}$ and the Goldberger-Treiman relation for N^* . When $g_A^{N^* N^*} = \pm 1$ for the naïve and mirror assignments, we find $g_{\pi N^* N^*} = g_A^{N^* N^*} m_{N^*}/f_\pi \sim \pm 17$. On the other hand, a recent work in the chiral unitary approach gives $g_{\pi N^* N^*} \sim 5$ ³⁵⁾. Here, just for simplicity, we use the same absolute value as $g_{\pi NN}$. The coupling values used in our computations are summarized in Table II.

Besides the resonance dominant contributions as shown in Fig. 6, there are several other possible terms. We ignore all of them from the following reasons.

- **Background:**



First we consider three diagrams as shown in Figs. 7 (a-c). All of them are not allowed due to G-parity. For (a), the vertex $\pi\pi\pi\eta$ is G-parity forbidden, since $G(\pi) = -1$ and $G(\eta) = 1$. For (b) and (c), to estimate the diagrams, we first consider the lowest order of the chiral expansion in the lagrangian³²⁾. For two-meson nucleon vertices in (b), the two mesons are correlated as vector mesons (such as ρ) which have G-parity plus. Therefore, the G-parity minus combination $\pi\eta$ is not allowed. Similarly, three meson vertices in (c) have axial vector correlation with negative G-parity, and hence the positive G-parity combination $\pi\pi\eta$ is not allowed. These selection rules are explicitly satisfied in chiral lagrangians.

- **ρ meson:**

We have computed the diagram in Fig. 7 (d) explicitly. The rho meson coupling to N and N^* is extracted from the helicity amplitudes $A_{1/2} \sim 0.08 \text{ GeV}^{-1/2}$ ¹¹⁾ using the vector meson dominance as shown in Fig. 7 (d). It turns out that the contribution to the cross section is negligibly small as compared to the resonance pole terms in Fig. 14 by about a factor 10^{-3} .

- **Other resonance contributions:**

Finally, one would expect contributions where two resonances appear in intermediate states as shown in Fig. 7 (e). Again we can ignore these diagrams safely, since there is no strong indication that baryon resonances couple to $N^*(1535)$ by emitting a pion¹¹⁾. In particular, the delta resonance which could be excited strongly by the incident pion couples only weakly to $N^*(1535)$, since the observed branching ratio of $N^*(1535) \rightarrow \Delta\pi$ is less than 10 %¹¹⁾. The contamination from resonances of isospin $I = 3/2$ may be estimated by $p\pi^+ \rightarrow p\pi^+\eta$.

Fig. 7. Various contributions to $\pi N \rightarrow \eta \pi N$.Fig. 8. Total cross section of $\pi N \rightarrow \eta N$. Data are taken from³⁶⁾.

Before discussing the process we are interested in, we mention that our present approach does not include initial state interaction which affects and reduce absolute values of cross sections by factor of a few times. We show in Fig. 8 the total cross section of $\pi + N \rightarrow \eta + N$, where theoretical cross sections computed using the present lagrangian and parameters as compared with experiments. Theoretical cross section is indeed smaller than experimental one by a factor nearly two at the peak position.

Now we compute the S -matrices. Reduction formula allows us to write

$$\begin{aligned}
 S_{fi} &= {}_{out}\langle p_f, k_\pi, k_\eta | k_i, p_i \rangle_{in} \\
 &= disc. + (iZ_\pi^{-1/2})^2 (iZ_\eta^{-1/2}) \int d^4x d^4y d^4z e^{ik_\pi x + ik_\eta y - ik_i z}
 \end{aligned}$$




Fig. 9. Definition of momentum variables in the center of mass (CM) frame.

$$(\square_x + m_\pi^2)(\square_y + m_\eta^2)(\square_z + m_\pi^2)_{out} \langle p_f | T(\pi^i(x)\eta(y)\pi^j(z)) | p_i \rangle_{in}. \quad (5.2)$$

The momentum variables are defined as shown in Fig. 9. In the perturbation theory, the T -matrix element can be computed as

$$\begin{aligned} &_{out} \langle p_f | T(\pi^i(x)\eta(y)\pi^j(z)) | p_i \rangle_{in} \\ &= i^3 \langle p_f | T(\pi^i(x)\eta(y)\pi^j(z)) \int d^4x_1 d^4x_2 d^4x_3 L_{\eta NN^*}(x_1) L_{\pi NN^*}(x_2) \\ &\quad \otimes (L_{\pi NN}(x_3) + L_{\pi N^* N^*}(x_3)) | p_i \rangle_{in}. \end{aligned} \quad (5.3)$$

After performing the Wick contraction in (5.3) which is then inserted in (5.2), we find an ordinary expression for amplitudes in momentum space. For instance the amplitudes for Fig. 14 (1) and (2) are given by

$$T(1) = \bar{u}(p_f) \frac{(i\sqrt{2}g_{\pi NN}i\gamma_5)i(i g_{\eta NN^*})i(i\sqrt{2}g_{\pi NN^*})}{(\not{p}_f + \not{k}_\pi - m_N)(\not{p}_i + \not{k}_i - m_{N^*} + \frac{i}{2}\Gamma)} u(p_i), \quad (5.4)$$

$$T(2) = \bar{u}(p_f) \frac{(i g_{\eta NN^*})i(i\sqrt{2}g_{\pi N^* N^*}i\gamma_5)i(i\sqrt{2}g_{\pi NN^*})}{(\not{p}_f + \not{k}_\eta - m_{N^*} + \frac{i}{2}\Gamma)(\not{p}_i + \not{k}_i - m_{N^*} + \frac{i}{2}\Gamma)} u(p_i), \quad (5.5)$$

where u 's are the Dirac spinors for the nucleon and $\Gamma = \Gamma_{N^*(1535) \rightarrow \pi N} + \Gamma_{N^*(1535) \rightarrow \eta N} = 140$ MeV is the total width of $N(1535)$.

Using the T -matrix, $\mathcal{T}_{fi} = T(1) + \dots T(6)$, we calculate differential cross section as

$$d\sigma = \frac{2m_N}{4\sqrt{(p_i \cdot k_i)^2 - m_N^2 m_\pi^2}} \frac{1}{2} \sum_{spin} |\mathcal{T}_{fi}|^2 d\Phi, \quad (5.6)$$

where the phase space of the three body final state is given by

$$d\Phi = (2\pi)^4 \delta(p_i + k_i - p_f - k_\pi - k_\eta) \frac{d^3k_\pi}{(2\pi)^3 2E_\pi} \frac{d^3k_\eta}{(2\pi)^3 2E_\eta} \frac{m_N d^3p_f}{(2\pi)^3 E_f}. \quad (5.7)$$

Here the convention for the normalization is

$$\bar{u}^{(\alpha)}(p) u^{(\beta)}(p) = \delta^{\alpha\beta}, \quad \langle p | p' \rangle = \frac{E}{m} (2\pi)^3 \delta^3(\vec{p} - \vec{p}'). \quad (5.8)$$

In the center of mass frame, the phase space integral measure reduces to¹¹⁾

$$d\Phi = \frac{m_N}{4(2\pi)^5} dE_\pi dE_f d\alpha d(\cos \beta) d\gamma. \quad (5.9)$$

Here the orientations of the momenta of the emitted particles are specified by the three Euler angles α , β and γ , since, in the center of mass frame, the three momenta, $\vec{p}_f$, $\vec{k}_\pi$ and $\vec{k}_\eta$, lie in a common plane. Then the relative angles between either two of $\vec{p}_f$, $\vec{k}_\pi$ and $\vec{k}_\eta$ can be determined, if the energies of the proton and the pion in the final state, E_f , E_π are fixed.

We have computed the integral over the three body phase space in the Monte Carlo method. The number of configurations is taken more than 30,000, depending on the kinds of cross section. The total cross section is computed in a schematic way as

$$\sigma = (\text{K.F.}) \int |\mathcal{T}(\xi)|^2 d\Phi \rightarrow (\text{K.F.}) \frac{1}{N} \sum_{i=1}^N |\mathcal{T}(\xi_i)|^2 V, \quad (5.10)$$

where $\xi = (E_f, E_\pi, \alpha, \cos \beta, \gamma)$. The volume of the phase space is given by the integral

$$V = \frac{m_N}{4(2\pi)^5} 4\pi^2 \int_{E_{min}}^{E_{max}} \frac{2\sqrt{(E_\eta^{*2} - m_\eta^2)(E_\pi^{*2} - m_\pi^2)}}{E_{cm}} dE_\pi, \quad (5.11)$$

where E_η^* and E_π^* are the energies of the emitted η and π in the rest frame of the emitted nucleon and eta. They are

$$E_\eta^* = \frac{E_{cm}^2 - m_N^2 + m_\eta^2 + m_\pi^2 - 2E_{cm}E_\pi}{2\sqrt{E_{cm}^2 + m_\pi^2 - 2E_{cm}E_\pi}}, \quad (5.12)$$

$$E_\pi^* = \frac{E_{cm}E_\pi - m_\pi^2}{\sqrt{E_{cm}^2 + m_\pi^2 - 2E_{cm}E_\pi}}. \quad (5.13)$$

The lower and upper bounds of the integral (5.11) are given by

$$E_{min} = m_\pi, \quad (5.14)$$

$$E_{max} = \frac{E_{cm}^2 + m_N^2 - (m_N + m_\eta)^2}{2E_{cm}}. \quad (5.15)$$

To compute differential cross section $d\sigma(\zeta)$ where ζ is a representative of the variable we need, for instance, the angle of the emitted pion in the center of mass frame and the momentum of the emitted pion in the laboratory frame. We insert a delta function of finite range $\Delta\zeta$ in the integrand of the total cross section.

$$\begin{aligned} d\sigma(\zeta) &= (\text{K.F.}) \int |\mathcal{T}(\xi)|^2 \delta(\zeta'(\xi) - \zeta) d\Phi \\ &\rightarrow (\text{K.F.}) \frac{1}{N} \sum_{i=1}^N |\mathcal{T}(\xi_i)|^2 \frac{1}{\sqrt{\pi}\Delta\zeta} e^{-\frac{(\zeta'(\xi_i) - \zeta)^2}{\Delta\zeta^2}} V. \end{aligned} \quad (5.16)$$




Fig. 10. (a) Total cross sections of $\pi^- p \rightarrow \eta \pi^- p$ for the naive and mirror models as functions of the initial pion momentum $P_{c.m.}$ in the center of mass frame. (b) Separate contributions from various terms of Fig. 14. The amplitude of the cross term $T(2)^* T(3)$ corresponds to the naive assignment. It changes the sign for the mirror assignment.



Note that the phase space is represented in the center of mass frame. The ζ' stands for the translation of the variables ξ . For example, ζ' is a boost transformation from the CM frame to the laboratory frame to calculate the differential cross section in the laboratory frame.

The total cross sections is shown in Fig. 10 (a) as functions of the energy of the initial pion for the naive and mirror cases. The difference between the two is due to the sign of the $g_{\pi N^* N^*}$ coupling. The cross sections increase, as the initial pion energy and correspondingly the phase space of the final three body state increase. For $P_{cm} \gtrsim P_{cm}^{\text{threshold}} + 50 \text{ MeV/c}$ ($P_{cm}^{\text{threshold}} = 528 \text{ MeV/c}$), the cross sections reach more than ten micro barn, which will be experimentally accessible. In the whole energy region as shown in Fig. 10 the cross section is larger in the mirror model, which is about twice as that in the naive model.

Among various terms shown in Fig. 14, major contributions are from the diagrams (2) and (3). In Fig. 10 (b), we show relative strengths of $|T(2)|^2$, $|T(3)|^2$ and their interference $2T(2)^* T(3)$ in the naive assignment. The difference between the naive and mirror assignments in the total cross section is from the sign difference of this term. The third major contribution is from $T(1)$ which is also shown there. Other terms are negligible.

In the pion induced reaction, the term $T(3)$ gives a large contribution to the cross section as compared with $T(1)$ and $T(2)$, although we expected naively that the terms $T(1)$ and $T(2)$ were the dominant contributions when considering their energy denominators. This can be explained by the p-wave nature of the $\pi N N$ and $\pi N^* N^*$ couplings. In the diagrams 1 and 2, a p-wave pion is emitted with a small momentum, which suppresses the amplitudes. In the diagram (3), a p-wave pion is absorbed with a large momentum, which enhances the amplitude.

In the present calculation, we have not considered the initial and final state interactions between the pion and nucleon. The inclusion of realistic interactions

Fig. 11. Angular distributions of the π^- in the final state.Fig. 12. Momentum distributions of the π^- in the final state.

might reduce absolute values of cross sections by an overall factor of several times, as we have discussed the single eta production $p\pi^- \rightarrow n\eta$ in Fig. 8. It is important to consider such distortion effect when we discuss absolute values of the cross sections. However, relative behaviors of such as momentum and angular distributions are not affected by this effect. These are now discussed as follows.

- Angular distributions:** In Fig. 11, we show angular distributions of the final state pion in the center of mass frame, which shows clear difference between the two models. This comes from the p-wave nature of the $\pi N^* N^*$ coupling in the diagram (2) of Figs. 14. As it is mentioned before, the main contributions are given by the diagram (2) and (3), and the difference in the sign of $g_{\pi N^* N^*}$ appears in the sign of the interference term of $T(2)$ and $T(3)$. Due to the structure of the Yukawa vertex, the emitted pion in the diagram (2) is in p-state, which gives linear dependence on $\cos\theta_{\pi}^{c.m.}$. On the other hand, the emitted pion in the diagram (3) is in s-state, which has no angular distributions. Therefore the cross term of $T(2)$ and $T(3)$ behaves linearly in $\cos\theta_{\pi}^{c.m.}$, and the apparently different behaviors in the $\cos\theta_{\pi}^{c.m.}$ dependence is due to the difference in the sign of the coupling $g_{\pi N^* N^*}$. This angular dependence would be one of the cleanest observables to distinguish the naive or mirror assignments.
- Momentum distributions:** Another example which is useful is the momen-




Fig. 13. A diagram of the Kroll-Ruderman type for a charged pion production.

tum (energy) distribution of one of the final state particles. We plot the momentum distribution of the emitted pion in the laboratory frame in Fig. 12 for several incident energies. What differs in the two chiral assignments is the position of the peak in the cross sections. In the naive case, it does not depend on the incident energy very much, while in the mirror case, it shifts to higher momentum region as the incident energy is increased.

5.3. Photon induced reaction: $\gamma + N \rightarrow \pi + \eta + N$

In general, photoproductions contain richer physics than pion induced processes, since the photon carries spin one and has isoscalar and isovector components. Our investigation is similar to what was explored for the two pion productions^{37), 38)}, but the process considered here is a little bit simpler since the eta meson is an isoscalar particle and exclusively couples to $N(1535)$. Under reasonable assumptions, we can follow very closely to the method for the pion induced reactions. In the following we discuss several items relevant to the photoproduction. Details will be reported elsewhere³⁹⁾.

First, as in the case of single pion photoproduction, we expect that the Kroll-Ruderman term as shown in Fig. 13 plays a dominant role for charged pion production. However, this term is suppressed for a neutral pion production. For this reason, in the following discussions we consider the neutral pion production: $\gamma p \rightarrow \pi^0 \eta p$.

Thus we compute the twelve resonance dominant diagrams as shown in Figs. 14 in which the initial pion is replaced by a photon. For γNN vertex, we have

$$\mathcal{L}_{\gamma NN} = -e\bar{N}\gamma_\mu \frac{1 + \tau_3}{2} N A^\mu + \frac{e}{4M}\bar{N}(\kappa_S + \kappa_V\tau_3)\sigma^{\mu\nu} N F_{\mu\nu} + \text{h.c.} \quad (5.17)$$

where the κ_S and κ_V are related to the isoscalar and isovector anomalous magnetic moments and take the values

$$\kappa_S = -0.12 \quad \kappa_V = 3.7. \quad (5.18)$$

For $\gamma N^* N^*$, we assume the same form as (5.17) but the nucleon mass is replaced by the resonance mass. The unknown magnetic moment of N^* ($\equiv \kappa^{**}$) is taken just the same as that of the nucleon. The uncertainty from this is, however, not very important, since the diagrams which contain the $\gamma N^* N^*$ coupling (Figs. 14 (5) and (8)) play only a minor role. For γNN^* vertex, we take the tensor form which is




Fig. 14. Twelve N^* dominant diagrams for $\gamma p \rightarrow \pi^0 \eta p$ process.

compatible with gauge invariance:

$$\mathcal{L}_{\gamma NN^*} = \frac{ie\kappa_V^*}{2(m_{N^*} + m_N)} \bar{N}^* \tau_3 \gamma_5 \sigma^{\mu\nu} N F_{\mu\nu} + \text{h.c.} \quad (5.19)$$

where we used isovector dominance and the magnetic coupling is given as

$$\kappa_V^* = 0.9, \quad (5.20)$$

which is determined from the analyses of eta photoproduction⁴⁰⁾.

We still need to consider signs of coupling constants carefully, since there are more coupling constants in the photon induced reaction than in the pion induced reaction. The twelve diagrams of Fig. 14 are classified into four groups according to the common coupling constants:

Group A $\sim$ Figs. (1) (4) (9) and (12) $\sim g_{\gamma NN^*} g_{\pi NN}$

Group B $\sim$ Figs. (2) and (11) $\sim g_{\gamma NN^*} g_{\pi N^* N^*}$

Group C $\sim$ Figs. (3) (6) (7) and (10) $\sim g_{\gamma NN} g_{\pi NN^*}$

Group D $\sim$ Figs. (5) and (8) $\sim g_{\gamma N^* N^*} g_{\pi NN^*}$

Here we do not include another coupling constant $g_{\eta NN^*}$ which is common to all the diagrams, since only relative signs are our concern. Also, the photon coupling

constants $g_{\gamma NN}$ etc are symbolic, since actual couplings contain both Dirac (vector) and Pauli (tensor) types as given in (5.17).

Since we are concerning only about relative signs, let us multiply $g_{\pi NN^*}$ to the above combinations:

$$(A) \rightarrow g_{\gamma NN^*} g_{\pi NN^*} g_{\pi NN} , \quad (5.21)$$

$$(B) \rightarrow g_{\gamma NN^*} g_{\pi NN^*} g_{\pi N^* N^*} , \quad (5.22)$$

$$(C) \rightarrow g_{\gamma NN} g_{\pi NN^*}^2 , \quad (5.23)$$

$$(D) \rightarrow g_{\gamma N^* N^*} g_{\pi NN^*}^2 . \quad (5.24)$$

The relative sign of $g_{\gamma NN^*}$ and $g_{\pi NN^*}$ is known from the pion photoproduction; it is plus for the proton and minus for the neutron⁴¹⁾. The opposite signs for the proton and neutron is due to isovector nature of the photon coupling to $N(1535)$. Since we know the photon coupling to the nucleon, the sign ambiguity among the groups (A), (B) and (C) is removed modulo the relative sign of $g_{\pi NN}$ and $g_{\pi N^* N^*}$, which is the one under the current interest. The sign of the last group (D) can not be determined; it depends on the sign of the coupling $g_{\gamma N^* N^*}$. The Dirac coupling is determined since it is the charge of $N(1535)$. However, the strength of the Pauli term can not be fixed; it depends on the unknown magnetic moments of $N(1535)$. It turns out, however, that the contribution of the two terms of the group D, Figs. 14 (5) and (8), are not very important. We have verified this by computing cross sections with different values of the anomalous magnetic moments of N^* . Typical cases are shown in Figs. 15 (see explanations below). In this way, we can again discuss the relative sign of $g_{\pi NN}$ and $g_{\pi N^* N^*}$ among the diagrams of the groups A, B and C.

As in the previous case, we have computed total cross sections as well as momentum and angular distributions. In Fig. 15, total cross sections are plotted as functions of the incident photon energy. Here we have shown the results using the anomalous magnetic moments of N^* which are the same as the nucleon and for those with opposite sign. The difference is within several ten percent, justifying relatively less importance due to the ambiguity in these unknown couplings. Unlike the pion induced reactions, the cross section of the naive model is larger than the mirror model. This is so because the dominant diagrams in the present process are (1) and (2) of Figs. 14 and the approximate cancellation between them are realized in the mirror model. Accordingly, the angular and momentum distributions shown in Figs. 16 follow this tendency in magnitude. The shape of momentum distributions are qualitatively the same for the naive and mirror models, while the angular distributions differ in the two assignments, reflecting once again the sign of the pion p-wave coupling.

§6. Conclusion

We have considered chiral symmetry aspects of baryons by assuming that they belong to suitable linear representations of the chiral symmetry group. We have emphasized a possibility that the positive and negative parity nucleons form a large representation of the chiral group where the positive and negative parity nucleons




Fig. 15. Total cross sections for $\gamma p \rightarrow \eta \pi^0 p$ for the naive and mirror models as functions of the initial pion momentum $P_{c.m.}$ in the center of mass frame.




Fig. 16. (a) Angular and (b) momentum distributions of π^0 in the final state. Cross sections at the incident photon energy $E = 590$ MeV are scaled by 10 for the angular distributions and 20/3 for the momentum distributions. Magnetic moments of N^* , κ^{**} , is taken to be the same as that of the nucleon.

transform to each other under axial transformations. This particular representation has been known for some time but we have shown that this is one of two possible representations when there are two nucleons.

A distinguished feature of the mirror representation is that the different parity nucleons can form a degenerate multiplet with a finite mass m_0 when chiral symmetry is restored. This is very much different from the ordinary linear sigma model, in which the nucleon masses are generated solely by spontaneous breaking of chiral symmetry. The origin of finite $m_0 \neq 0$ is beyond the dynamics of chiral symmetry. It is interesting if more information can be extracted from lattice QCD or QCD sum rules methods.

Another aspect of the mirror model is that the negative parity nucleon carries the negative axial charge. In effective theories, it is a group theoretical consequence.

However, microscopic origin should be investigated, as baryons are composite objects. Further study of this is absolutely necessary³⁵⁾.

After establishing general features of the chiral symmetry representations for baryons, we have studied microscopic origin by using baryon interpolating fields. It turns out that widely used interpolating fields of three quarks possess the naive chiral property. This can be seen both directly through commutation relations and indirectly by computing the Yukawa pion coupling. By contrast, the mirror chiral property is realized by five quark operators. This can be shown once again both in commutation relations and pion couplings.

Finally, we have considered experiments to observe the theoretical issues discussed here. There is a fortunate situation that the negative parity resonance $N(1535)$ can be seen almost selectively by detecting eta in the final state. Using this, we have proposed eta and pion productions in pion or photon induced reactions. interference effects are then used to observe the relative sign of $g_{\pi NN}$ and $g_{\pi N^* N^*}$ which can distinguish either naive or mirror property of the nucleon.

Although chiral symmetry has a long history in strong interaction physics, its realization in the baryon sector has not been investigated very much so far. It serves an interesting subject in future hadron physics.

Acknowledgements

We would like to thank Prof. T. Kunihiro, Prof. T. Hatsuda, Dr. H. Kim and Dr. Y. Nemoto for discussions. This work is supported in part by the Grant-in-Aid for scientific research (C)(2) 11640261. The work by D.J. is supported by the Spanish Ministry of Education in the Program “Estancias de Doctores y Tecnólogos Extranjeros en España”.

References

- 1) Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961), 345.
- 2) J. Goldstone, Nuovo Cim. **19** (1961), 154.
- 3) S. Weinberg, *Quantum Theory of Fields*, Cambridge Univ. Press, Cambridge (1985).
- 4) S. Weinberg, Phys. Rev. **177** (1969), 2604.
- 5) I.S. Gerstein and B.W. Lee, Phys. Rev. Lett. **16** (1966), 1060.
- 6) S. Weinberg, Phys. Rev. **166** (1968), 1568.
- 7) J. Gasser and H. Leutwyler, Phys. Rept. **87** (1982) 77.
- 8) G. Ecker, Prog. Part. Nucl. Phys. **35** (1995) 1.
- 9) A. Pich, Rept. Prog. Phys. **58** (1995) 563.
- 10) M. Gell-Mann and M. Levy, Nuovo Cim. **16** (1960) 705.
- 11) Particle Data Group, Euro. Phys. J. **51** (2000) 1.
- 12) D. Jido and M. Oka and A. Hosaka, Phys. Rev. Lett. **80** (1998) 448.
- 13) Y. Nemoto, D. Jido, M. Oka and A. Hosaka, Phys. Rev. **D57** (1998) 4124.
- 14) D. Jido, Y. Nemoto, M. Oka and A. Hosaka, Nucl. Phys. A671 (2000) 471.
- 15) D. Jido, M. Oka and A. Hosaka, preprint (2000) hep-ph/0007127.
- 16) D. Jido, T. Hatsuda and T. Kunihiro, Phys. Rev. Lett. **84**, 3252 (2000).
- 17) H. Kim, D. Jido and M. Oka, Nucl. Phys. **A640**, 77 (1998).
- 18) B. W. Lee, *Chiral Dynamics*, Gordon and Breach, New York, (1972).
- 19) C. DeTar and T. Kunihiro, Phys. Rev. **D39** (1989) 2805.
- 20) T.D. Cohen and X. Ji, Phys. Rev. **D55** (1997), 6870.
- 21) T. Hatsuda and M. Prakash Phys. Lett. **B224** (1989) 11.

- 22) S.R. Beane, Phys. Lett. B444, 147, (1998).
- 23) G.A. Christos, Phys. Rev. D35, 330, (1987); Z. Phys. C29, 361, (1985); *ibid.* C21, 83, (1983).
- 24) J. Wess and B. Zumino, Phys. Lett. **B37**, 95 (1971); E. Witten, Nucl. Phys. **B223**, 442 (1983).
- 25) R. Koniuk and N. Isgur, Phys. Rev. **D21** (1980), 1868.
- 26) R. Bijker, F. Iachello and A. Leviatan, Ann. Phys. **236** (1994) 69; Phys. Rev. **D55** (1997), 2862.
- 27) H. Yamazaki et. al., Nucl. Phys. **A670**, 202 (2000).
- 28) T. Yorita et. al., Phys. Lett. **B476**, 226 (2000).
- 29) D. Jido, N. Kodama and M. Oka, *Phys. Rev. D* **54**, 4532 (1996); D. Jido and M. Oka, hep-ph/9611322.
- 30) B. L. Ioffe, Nucl. Phys. **B188**, 317 (1981).
- 31) D. Espriu and P. Pascual and R. Tarrach, Nucl. Phys. **B214**, 285 (1983).
- 32) See, for instance, J.F. Donoghue, E. Golowich and B.R. Holstein, *Dynamics of the standard model*, Cambridge monographs on particle physics (1992).
- 33) D.M. Manley and E.M. Salesky, *Phys. Rev. D* **45**, 4002 (1992).
- 34) T.P. Vrana, S/A. Dytman and T.-S.H. Lee, Phys. Reports 328, 181 (2000).
- 35) J.C. Nacher, A. Parreño, E. Oset A. Ramos, A. Hosaka and M. Oka, Nucl. Phys. **A678**, 187 (2000).
- 36) M. Batinic, I. Slaus, A. Svarc and B. M. K. Nefkens, Phys. Rev. **C51** 2310, (1995).
- 37) K. Ochi, M. Hirata and T. Takaki, Phys. Rev. **D56** 1472, (1997).
- 38) J.C. Nacher, E. Oset, M.J. Vicente Vacas, L. Roca, nucl-th/0012065, to appear in Nucl. Phys. A.
- 39) D. Jido, M. Oka and A. Hosaka, in preparation.
- 40) M. Benmerrouche, Nimai C. Mukhopadhyay and J. F. Zhang, Phys. Rev. **D51** 3237 (1995).
- 41) Particle Data Group, Rev. Mod. Phys., **48** Part II, 1 (1976).